

The place of CEUS in distinguishing benign from malignant cervical lymph nodes: a prospective study

Laura Poanta¹, Oana Serban¹, Isabela Pascu¹, Sever Pop², Marcel Cosgarea², Daniela Fodor¹

¹2nd Internal Medicine Department, ²ENT Department, "Iuliu Hatieganu" University of Medicine and Pharmacy, Cluj-Napoca, Romania

Abstract

Aims: The aim of the study was to evaluate the ability of contrast-enhanced ultrasonography (CEUS) compared to gray-scale B-mode and color Doppler ultrasound in differentiate benign versus malign superficial cervical lymph nodes. **Material and methods:** In a prospective study ultrasonography (gray scale, color and spectral Doppler, and CEUS) was performed in 61 patients (33 men, 28 women; mean age of 51.2 years, range: 18-81 years), with cervical lymphadenopathy. The nodes were examined and biopsied or surgically removed. CEUS was performed with 2.4 ml intravenous bolus of contrast agent Sono Vue and the results were registered with a special software. **Results:** Of all the nodes, 32 were benign and 29 were malignant (metastases). Solbiati index was higher in benign nodes (2.23 ± 0.84 vs 1.50 ± 0.48 , $p < 0.05$). Doppler parameters (vessel location, vascular pattern, pedicullum number, resistivity index, and pulsatility index) were significantly lower in benign nodes ($p < 0.001$), and ROC analysis returned excellent results. For CEUS, derived peak intensity (DPI %) was higher in benign nodes (17.72 ± 5.43 vs 11.76 ± 4.88 , $p < 0.05$); regional blood volume (RBV) was also higher (849.8 ± 467.1 vs 458.3 ± 283.3 , $p < 0.05$). The time to peak (TTP, s) and area under the curve (AUC, cm²) were similar in both benign and malignant nodes. Enhancement pattern was the most accurate to characterize benign versus malignant nodes. Sensitivity and specificity were higher for DPI, RBV and enhancement pattern from CEUS, according to ROC analysis, compared to gray scale ultrasound, but lower than color Doppler. Analyzing the place of CEUS in lymph node evaluation we found that CEUS is most useful for the evaluation of the lymph nodes with uncertain aspect at gray scale and Doppler evaluation. **Conclusions:** ROC analysis confirmed the higher degree of diagnostic accuracy of CEUS in comparison with conventional techniques for some parameters such as enhancement pattern. Evaluation of nodal perfusion with this method can be helpful in the differentiation of benign from malignant nodes but requires further confirmation.

Keywords: lymph nodes, ultrasonography, gray scale, Doppler, CEUS

Introduction

Ultrasound (US) is the commonly used method for evaluating superficial lymph nodes and it is also important in staging nodal metastases [1,2]. It is widely available, easy to use, and less expensive than other imaging

methods, and it has the benefit of nonionizing radiation. Accurate lymph node characterization is important for a wide number of clinical situations, including prognosis prediction, selecting and monitoring treatment, beyond the diagnosis itself (cancer, lymphoma, or inflammatory nodes) [1,3]. The differences between benign and malignant nodes are reported on the basis of various diagnostic criteria in gray scale US, such as the shape, border, echogenicity, calcification, necrosis, or vascularization pattern by Doppler US examination. Malignant nodes are described to be hypoechoic, without echogenic hilum, with round shape, longitudinal to transverse diameters ratio less than 2, and blood vessels predominant in periphery [3-7]. Contrast-enhanced ultrasound (CEUS) is a modern imaging method evaluating tissue perfusion in real time. The administration of contrast agent intrave-

Received 16.12.2013 Accepted 18.01.2014

Med Ultrason

2014, Vol. 16, No 1, 7-14

Corresponding author: Daniela Fodor, MD, PhD

2nd Internal Medicine Department

"Iuliu Hatieganu" University of Medicine and

Pharmacy, Cluj-Napoca, Romania

2-4 Clinicilor str

400006 Cluj Napoca, Romania

Phone: 004 0264591942/442

Email: dfodor@umfcluj.ro

nously and special soft of ultrasound device is necessary [8,9]. In our prospective study we analyzed the usefulness of CEUS in distinguishing between benign and malignant lymph nodes, evaluating the nodes perfusion and enhancement pattern compared with conventional US, in the differential diagnosis of the superficial lymphadenopathy. The results were compared with the histological examination of the nodes.

Material and methods

Prospectively, 61 patients (33 men, 28 women; mean age of 51.2 years, range: 18-81 years) were examined by US (gray scale, Doppler, and CEUS) between September 2012-March 2013 in the 2nd Internal Department. The majority of the patients were referred for US examination from ENT department due to the existence of palpable mass/masses in the cervical region. For this reason all the malignant nodes represented metastasis from ENT cancers. In cases with multiple lymphadenomegaly the most representative lymph node was chosen for analysis. The diagnoses were confirmed by US guided biopsy or by surgical removal of the lymph node and subsequent histological examination. Pediatric patients and patients with lymphoma were excluded from the study. The local Ethics Committee approved the study, and all participants gave their written informed consent prior to the investigation.

Gray scale and color Doppler US examination

All the patients were examined with a Logiq S8 GE ultrasound machine with a high-frequency linear probe (6–15MHz).

In gray scale the following parameters were recorded: the longitudinal and transverse diameters (for Solbiati index calculation - the ratio between the longitudinal and the transversal axis), the echogenicity of the nodes (classified as hypoechoic, isoechoic, and hyperechoic), hilum visibility (present, hard to see, or absent), margins (sharp, irregular, blurred), and homogeneity or internal structure changes (calcification, necrosis).

In Doppler US (pulse repetition frequency 350 Hz, wall filter 45 Hz) three patterns of nodal vascularization were defined: 1) hilar pattern with flow signals in the nodal hilum, 2) peripheral pattern with flow signals mainly in peripheral nodal parts, and 3) mixed pattern with both hilar and peripheral pattern. The resistivity index (RI) and pulsatility index (PI) were measured in the main artery of the node.

CEUS examination

For CEUS examination (3–9 MHz linear probe, range of gain: 80–94%, compression 36, mechanical index 0.07) a bolus of 2.4 ml of contrast agent (Sono-




Fig 1. CEUS examination, peak of signal intensity map, and CEUS parameters of an inflammatory lymph node.




Fig 2. CEUS examination, peak of signal intensity map, and CEUS parameters of a metastasis from laryngeal cancer.

Vue, Bracco Imaging SpA, Milan, Italy) was administered intravenously, followed by the flushing of 10 ml saline solution. All CEUS examinations were digitally recorded. The nodal perfusion and the enhancement pattern were evaluated in the arterial (10–15 s after bolus of contrast agent) and parenchymal (15–30 s after bolus of contrast agent) phase. Four nodal enhancement patterns were defined: 1) intense homogeneous enhancement, 2) moderate homogeneous enhancement, 3) inhomogeneous enhancement with perfusion defects, and 4) lack of enhancement. Using Qontrast software (Esaote S.p.a, Florence, Italy) for perfusion quantification the derived peak intensity (DPI - %), time to peak (TTP- seconds), regional blood volume (RBV) and area under the curve (AUC – cm²) were measured (fig 1, fig 2).

After the histological examination of the biopsy material or of the surgical removed node, the examined cervical lymph nodes were divided into two groups: benign and malignant nodes and the US findings were compared with histological diagnosis.

Statistical analysis

Standard descriptive statistics were adopted for the analysis; we defined means and standard deviations for numeric results. T-test was used to compare means. Chi-square was used to compare the categorical parameters. ROC (Receiver Operating Characteristic) analysis was

Table I. Lymph nodes characterization by gray scale ultrasound.

Ultrasound characteristic n(%)	Benign nodes (N=32)	Malignant nodes (N=29)	P value
SI (L/T \geq 2)	18 (56.2)	2 (6.8)	<0.001
Echogenicity			
Hypoechoic	8 (25)	18 (62)	
Isoechoic	17 (53.1)	9 (31)	0.011
Hyperechoic	7 (21.8)	2 (6.8)	
Hilum visibility			
Present	24 (75)	11 (37.9)	
Hard to see	8 (25)	15 (51.7)	<0.001
Absent	0	3 (10.3)	
Internal structure			
Homogenous	25 (78.1)	6 (20.6)	
Inhomogeneous	7 (21.8)	21 (72.4)	<0.001
Necrosis	0	0	
Calcifications	0	2 (6.8)	
Margins			
Sharp	26 (81.2)	15 (51.7)	
Irregular	0	5 (17.3)	0.009
Blurred	6 (18.8)	9 (31)	

Legend: SI = Solbiati index. L/T = ratio between longitudinal and transverse diameters

used for the evaluation of sensitivity, specificity, positive and negative predictive value and overall ability of contrast-enhanced ultrasonography, gray-scale B-mode and power Doppler parameters to distinguish between benign and malignant nodes by calculating the area under the curve (we named it AUROC not to be confused with CEUS software measured parameter AUC). P value was considered significant at a value lower than 0.05. The results among all the investigation techniques were compared in order to establish a possible examination protocol and the place of CEUS in lymph nodes US investigation. Microsoft Office Excel 2003 and SPSS for Windows version 19 was used.

Results

Gray scale examination

The gray scale characteristics of the lymph nodes are detailed in table I. Mean L/T ratio was higher in the benign nodes ($p < 0.001$). Most of the benign nodes were iso- or hyperechoic (total 24 nodes, over 70%), while 62% of the malignant nodes were hypoechoic. Hilum was visible in 75% of the benign nodes, but in only 37.9% of the malignant nodes. Most of the benign nodes were homog-

enous, without necrosis or calcifications, and with sharp margins. Most of the malignant nodes were inhomogeneous, a minority had calcifications, and almost half of them had irregular or blurred margins.

Color Doppler examination

Color Doppler US showed different patterns in malignant versus benign nodes, (table II). Also, the mean values for RI and PI were lower in benign lymph nodes: 0.53 ± 0.06 in benign nodes vs 0.69 ± 0.07 in malignant nodes ($p < 0.001$), and 0.82 ± 0.16 in benign nodes vs 1.37 ± 0.35 in malignant nodes, respectively ($p < 0.001$). All of the benign nodes had hilar and regular vascular pattern, with only one pedicellum (31 of 32 nodes). More than half of the malignant nodes had peripheral or mixed vessels with chaotic patterns, and multiple pedicellus.

CEUS examination

CEUS enhancement patterns and perfusion quantification are shown in table III. DPI and RBV are significantly higher in benign nodes, while AUC and TTP showed similar results. The majority of the benign nodes showed homogenous enhancement patterns, while the malignant nodes showed mostly inhomogeneous patterns and even one node with no enhancement.

ROC analysis was performed for numerical values (table IV, fig 3 and 4).

To find the utility of the three US methods for correct identification of benign vs malignant nodes, the results were combined. Parameters used for analysis were the parameters found to have statistically significance in comparison benign vs malignant: for gray scale- L/T ratio, echogenicity, homogeneity, margins and hilum; for Doppler- vessel location, vascular pattern, pedicullum number, RI (cutoff value of 0.605 obtained from ROC analysis) and PI (cutoff value of 0.995 obtained from ROC analysis); and for CEUS- enhancement pattern, PI (cutoff value of 14.15 obtained from ROC analysis) and RBV (cutoff value of 497 obtained from ROC analysis).

If the gray scale US was combined with Doppler, all benign lymph nodes were correctly identified (100%), but only 22 from 29 malignant lymph nodes (75.8%). Six out of 29 malignant nodes were uncertain (20.7%) and one lymph node was incorrectly identified as benign (3.5%). When CEUS was added, five out of the six uncertain lymph nodes were correctly identified. It means that if all the three methods were combined 100% of benign nodes were correctly identified and 93.1% (27 from 29) of malignant nodes.

Considering the aforementioned parameters we developed a score that could be used to distinguish between malignant and benign lymph nodes using gray scale, Doppler and CEUS (table V).

Table II. Lymph nodes characterization by color Doppler – vascular pattern.

Nodes	Hilar pattern	Peripheral pattern	Mixed pattern	Regular pattern	Chaotic pattern	One pedicullum	Multiple pedicullus
Benign (N=32)	32 (100)	0	0	32 (100)	0	31 (96.8)	1 (3.1)
Malignant (N=29)	13 (44.8)	7 (24.1)	9 (31)	9 (31)	20 (68.9)	12 (41.4)	17 (58.6)
P value		<0.001			<0.001		<0.001

The results are expressed in number (%)

Table III. Lymph nodes characterization by CEUS – enhancement homogeneity and the visibility of hilar artery.

CEUS characteristic	Benign nodes (N=32)	Malignant nodes (N=29)	p value
Intense homogeneous	20 (62.5)	4 (13.8)	
Mild homogeneous	9 (28.1)	5 (17.2)	<0.001
Inhomogeneous	3 (9.3)	19 (65.5)	
No enhancement	0	1 (3.4)	
DPI (%)	17.72 ± 5.43	11.76 ± 4.88	<0.001
TTP (sec)	16.16 ± 2.90	17.12 ± 2.45	0.189
AUC (cm2)	1.11 ± 0.61	1.83 ± 1.73	0.036
RBV	849.8 ± 467.1	458.3 ± 283.3	<0.05

The results are expressed in number (%) or in number ± standard deviation. **Legend:** DPI = derived peak intensity, TTP = time to peak, AUC = area under the curve, RBV = regional blood volume

Table IV. ROC analysis: benign versus malignant nodes.

Parameters	AUROC value	P value	Statistical Sign**	CI 95%	Cutoff value	Sensitivity (%)	Specificity (%)
L/T ratio*	0.745	0.001	Fair	0.612 – 0.878	2*	93	53
RI	0.945	0.000	Excellent	0.890 – 0.999	0.605	90.3	84.4
PI	0.927	0.000	Excellent	0.865 – 0.990	0.995	87.1	84.2
RBV	0.746	0.001	Fair	0.620 – 0.872	497	75.9	75
DPI (%)	0.805	0.000	Good	0.696 – 0.914	14.15	75.9	75

Legend: *for L/T the cutoff value is pre-established, ** according to the following classification: 0.500 – 0.600 = fail, 0.600 – 0.700 = poor, 0.700 – 0.800 = fair, 0.800 – 0.900 = good, 0.900 – 1 = excellent. L/T ratio = ratio between longitudinal and transverse diameters, RI = resistivity index, PI = pulsatility index, RBV = regional blood volume, DPI = derived peak intensity.




Fig 3. ROC analysis for color Doppler (RI, resistivity index and PI, pulsatility index)




Fig 4. ROC analysis for gray scale and CEUS (DPI, derived pulsatility index, RBV, regional blood volume)

Table V. The score for distinguishing between malignant and benign nodes using gray scale US, color Doppler US, and CEUS.

Criteria	Characteristics	Number of Points
Gray scale US		
1. L/T ratio	≥ 2	0
	< 2	1
2. Echogenicity	Isoechoic/Hyperechoic	0
	Hypoechoic	1
3. Internal structure	Homogenous	0
	Inhomogeneous/Necrosis/ Calcifications	1
4. Margins	Sharp	0
	Irregular/Blurred	1
5. Hilum	Present	0
	Hard to see/Absent	1
Color Doppler US		
6. Vessel location	Hilar pattern	0
	Peripheral pattern/Mixed pattern	1
7. Vascular pattern	Regular pattern	0
	Chaotic pattern	1
8. Pedicullum number	One	0
	Multiple	1
9. RI	< 0.605	0
	≥ 0.605	1
10. PI	< 0.995	0
	≥ 0.995	1
Score < 5 – benign node		Score = 5 – CEUS needed
		Score > 5 – malignant node
CEUS		
Enhancement pattern	Homogenous	0
	Inhomogeneous/ No enhancement	1
DPI	≥ 14.15	0
	< 14.15	1
RBV	≥ 497	0
	< 497	1
Score ≤ 1 – benign node		Score ≥ 2 – malignant node

Legend: RI = resistivity index, PI = pulsatility index DPI = derived peak intensity, RBV = regional blood volume

If, after gray scale US and Doppler examination, less than 5 criteria are accomplished (<5 points), the nodes should be interpreted as benign. If more than 5 criteria are met (>5 points), the node should be interpreted as malignant. When the node accomplishes 5 criteria (5 points), it should be interpreted as uncertain and CEUS should be performed. After CEUS, if the node meets at least 2 criteria (≥ 2 points) it should be considered malignant.

Discussions

The role of US in the evaluation of superficial lymph nodes is well known and established [4,5]. It allows a detailed evaluation of nodes, including nodal internal structure and it seems to be superior to computed tomography (CT) for superficial lymph nodes [10]. The main utility of CT is the detection of deeply located nodes, especially in cervical region [10]. However, the limits of US are often discussed, such as incapacity to detect blood flow in all intranodal blood vessels by color and power Doppler [11,12].

Yu et al [13] studied 94 enlarged superficial lymph nodes. Regarding gray scale ultrasound, they showed that an L/T ratio of 2 or less has low specificity and accuracy. They found that L/T ratio is one of the less valuable parameters in the evaluation of lymph nodes. In our study, at a cut off value of 2, we have a very high sensitivity (93%) but a very low specificity (53%). Cutoff level of L/T ratio is 2 in most of the studies using gray scale US and it is already accepted that nodes with LT ratio ≥ 2 are classified as benign, while nodes with LT ratio <2 are malignant [14]. As for echogenicity, hypoechoic nodes are generally considered to be malignant. Also, absent hilum is suggestive for malignancy [7,14]. But it is important to note that in half of the malignant nodes in our study hilum was hard to see, so the results may be biased.

AUROC for L/T ratio, echogenicity, and hilum visibility returned values between 0.700 and 0.800, which, according to statistical significance, indicate a fair accuracy in distinguishing between the two groups of lymph nodes. So, gray B mode US is fairly reliable in describing lymph nodes, but has some limitations; it seems that grey scale US can identify malignant nodes but cannot correctly identify the lymph nodes as non-malignant. Other studies showed that CEUS improves the specificity in diagnosing benign lymph nodes as compared to B-mode US. It does not improve the correct identification of malignant lymph nodes and cannot replace EUS-guided fine-needle aspiration [15].

In our study, all the benign nodes showed hilar vascularity with a regular pattern. RI and PI were significantly

higher in malignant nodes and ROC analysis showed excellent results, which mean that this method could be considered as an excellent technique for differentiating benign from malignant nodes.

Currently used contrast agents allow a better description of microvascular pattern and can detect avascular areas of necrosis and tumoral metastasis [16]. In malignant nodes, neo-vessels with pathological aberrant feeding vessels are present, the growth being provided by angiogenic factors.

On the basis of the presence or absence of tumor angiogenesis there are various enhancement patterns of benign and malignant nodes described in literature [16-18]: benign nodes as intense homogeneous with no perfusion defects and with visible hilum artery, probably due to rich cortical capillary vascularity [19] or inhomogeneous enhancement and perfusion defects, with no visible artery in hilum, for malignant nodes [16-18].

Rubaltelli et al. [20] investigated the lymph nodes with focal cortical thickening. They found perfusion defects of thickened cortex in nodes affected by metastasis of cutaneous melanoma. They documented high specificity of CEUS, proved by histopathology. In our study the sensitivity and specificity were not as high as the Rubaltelli et al study, but we have similar results for enhancement patterns. We found the majority of benign lymph nodes to have homogenous enhancement on CEUS, either intense or moderate (29 of 32 nodes) and only three showed inhomogeneous enhancement. According to many authors [8,9,20] normal and reactive lymph nodes have intense vascularization with cortical capillary circulation and this fact could explain the homogenous aspect. On the other hand 19 (65.5%) of malignant nodes in our study, were inhomogeneous, and one of them had no enhancement, probably due to the fact that they were less vascularized and had perfusion defects [20]. CEUS is clinically valuable for microvascular analysis as it can deliver better data about vascular pattern and vascular defects. But few reports about the use of CEUS in superficial lymphadenopathy have been published and if they were, with controversial results [13].

ROC analysis for DPI and RBV returned good and fair results, respectively, but the values were significantly higher in benign nodes. According to some studies, TTP and AUC are considered to be lower in malignant nodes, but for DPI and RBV literature data are still contradictory [1,20,21].

Other studies [22] showed no differences between CEUS parameters (DPI, TTP and AUC) in benign versus malignant nodes, but only for the difference between maximum and minimum signal intensity (SI_{max} - SI_{min}). According to Quifang et al [22] CEUS patterns

differ considerably in benign versus malignant as nodes with metastases are vascularized with a heterogeneous centripetal enhancement pattern, and benign nodes have a homogenous centrifugal enhancement pattern. They underlined that even in metastatic nodes blood flow may be decreased due to vascular compression by neoplastic tissue, or vascular encasement. The blood in metastatic nodes may be affected also by encasement of the surrounding parenchyma [22]. This may explain the fact that peak intensity (DPI) varies widely among studies.

In our study, DPI and RBV were significantly lower in malignant nodes, while TTP and AUC were higher, but the difference between the two groups was statistically significant only for AUC. According to ROC analysis, these last two parameters are not useful in distinguishing between benign and malignant nodes, but AUROC had a good value for DPI and a fair value for RBV. The fact that TTP and AUC were higher in our study may be related to the technique we chose (the selection of the whole lymph node instead of a limited region of interest - ROI).

We established the criteria we used for malignancy from existing literature [8,9,23,24] and we also established cutoff values for RI, PI, DPI and RBV, from our ROC analysis. When we compared the results from all three techniques used in our study, we demonstrated that the correct diagnosis of uncertain nodes after grey scale and Doppler US was significantly increased by adding CEUS. This conclusion may be biased by the small number of patients. The main conclusion drawn from this study is that CEUS should be not performed as a routine examination, but only in these cases where the results are uncertain after using both gray scale and Doppler US.

The technique used in CEUS evaluation is very important, and the results may be influenced by the contrast agent used, patient characteristics and metabolism, and the selection of ROI. A limitation of this study, beside the small number of patients, is the lack of comparison with lymphoma lymph nodes.

Conclusions

In our study we identified different enhancement patterns in benign and malignant cervical lymph nodes. Our results showed a higher degree of diagnostic accuracy of CEUS in comparison with gray scale US, although the results obtained from ROC curves are not excellent. We did not demonstrate the superiority of CEUS over color Doppler for the assessment of nodal perfusion but CEUS can be helpful in the evaluation of nodal architecture and in the detection of abnormal vascular patterns, especially where the results are not certain. We found that Doppler US (color and spectral) is the best method for differen-

tiation in the benign versus malignant nodes, but more studies are necessary to establish the best ROI, the cutoff values for CEUS parameters, and to describe more accurately the enhancement patterns.

Conflict of interest: none

References

1. Slaisova R, Benda K, Jarkovsky J, Petrasova H, Szturz P, Valek V. Contrast-enhanced ultrasonography compared to gray-scale and power doppler in the diagnosis of peripheral lymphadenopathy. *Eur J Radiol* 2013; 82: 693-698.
2. Vassallo P, Wernecke K, Roos N, Peters PE. Differentiation of benign from malignant superficial lymphadenopathy: the role of high-resolution US. *Radiology* 1992; 183: 215-220.
3. Van den Brekel MW, Stel HV, Castelijns JA, et al. Cervical lymph node metastasis: assessment of radiologic criteria. *Radiology* 1990; 177: 379-384.
4. Ying M, Ahuja A. Sonography of neck lymph nodes. Part I: normal lymph nodes. *Clin Radiol* 2003; 58: 351-358.
5. Ahuja A, Ying M. Sonography of neck lymph nodes. Part II: abnormal lymph nodes. *Clin Radiol* 2003; 58: 359-366.
6. Solbiati L, Cioffi V, Ballarati E. Ultrasonography of the neck. *Radiol Clin North Am* 1992; 30: 941-954.
7. Dudea SM, Lenghel M, Botar-Jid C, Vasilescu D, Duma M. Ultrasonography of superficial lymph nodes: benign vs. malignant. *Med Ultrason* 2012; 4: 294-306.
8. Ahuja A, Ying M, HO SY, et al. Ultrasound of malignant cervical lymph nodes. *Cancer Imaging* 2008; 8: 48-56.
9. Cui XW, Jenssen C, Saftoiu A, Igonee A, Dietrich CF. New ultrasound techniques for lymph node evaluation. *World J Gastroenterol* 2013; 19: 4850-4860.
10. Rubaltelli L, Proto E, Salmaso R, Bortoletto P, Candiani F, Cagol P. Sonography of abnormal lymph nodes in vitro: correlation of sonographic and histologic findings. *AJR Am J Roentgenol* 1990; 155: 1241-1244.
11. Tschammler A, Ott G, Schang T, Seelbach-Goebel B, Schwager K, Hahn D. Lymphadenopathy: differentiation of benign from malignant disease—color Doppler US assessment of intranodal angioarchitecture. *Radiology* 1998; 208: 117-123.
12. Arijji Y, Kimura Y, Hayashi N, et al. Power Doppler sonography of cervical lymph nodes in patients with head and neck cancer. *AJNR Am J Neuroradiol* 1998; 19: 303-307.
13. Yu M, Liu Q, Song HP, et al. Clinical application of contrast-enhanced ultrasonography in diagnosis of superficial lymphadenopathy. *J Ultrasound Med* 2010; 29: 735-740.
14. Wunderbaldinger P. Problems and prospects of modern lymph node imaging. *Eur J Radiol* 2006; 58: 325-337.
15. Hocke M, Menges M, Topalidis T, Dietrich CF, Stallmach A. Contrast-enhanced endoscopic ultrasound in discrimination between benign and malignant mediastinal and abdominal lymph nodes. *J Cancer Res Clin Oncol* 2008; 134: 473-480.

16. Solbiati L, Cova L. Improved characterization of reactive and malignant lymph nodes using contrast-enhanced ultrasound. In: Albrecht T, Thorelius L, Solbiati L, Cova L, Frauscher F. (eds). *Contrast-enhanced ultrasound in clinical practice. Liver, prostate, pancreas, kidney and lymph nodes.* Springer-Verlag, Milan 2005: 39-50.
17. Döme B, Hendrix MJ, Paku S, Tóvári J, Tímár J. Alternative vascularization mechanisms in cancer: Pathology and therapeutic implications. *Am J Pathol* 2007; 170: 1-15.
18. Rubaltelli L, Khadivi Y, Tregnaghi A, et al. Evaluation of lymph node perfusion using continuous mode harmonic ultrasonography with a second-generation contrast agent. *J Ultrasound Med* 2004; 23: 829-836.
19. Gadre A, Briner W, O'Leary M. A scanning electron microscope study of the human cervical lymph node. *Acta Otolaryngol* 1994; 114: 87-90.
20. Rubaltelli L, Beltrame V, Tregnaghi A, Scagliori E, Frigo AC, Stramare R. Contrast-enhanced ultrasound for characterizing lymph nodes with focal cortical thickening in patients with cutaneous melanoma. *AJR Am J Roentgenol* 2011; 196: W8-W12.
21. Wilson SR, Greenbaum LD, Goldberg BB. Contrast-enhanced ultrasound: what is the evidence and what are the obstacles? *AJR Am J roentgenol* 2009; 193: 55-60.
22. Ouyang Q, Chen L, Zhao H, Xu R, Lin Q. Detecting metastasis of lymph nodes and predicting aggressiveness in patients with breast carcinomas. *J Ultrasound Med* 2010; 29: 343-352.
23. Na GD, Lim HK, Byun HS, Kim HD, Ko YH, Baek JH. Differential diagnosis of cervical lymphadenopathy: usefulness of color Doppler sonography. *AJR Am J Roentgenol* 1997; 168: 1311-1316.
24. Ouyang Q, Chen L, Zhao H, Xu R, Lin Q. Detecting metastasis of lymph nodes and predicting aggressiveness in patients with breast carcinomas. *J Ultrasound Med* 2010; 29: 343-352.